

POSTER PRESENTATION

Open Access

Efficient estimation of phase response curves via compressive sensing

Sungho Hong^{1*}, Erik De Schutter^{1,2}

From Twentieth Annual Computational Neuroscience Meeting: CNS*2011
Stockholm, Sweden. 23-28 July 2011

The phase-response curve (PRC), relating the phase shift of the oscillator and externally given perturbation [1], is one of the most important tools for characterizing the biological oscillators, such as neurons, and studying their synchronization in a network [2]. In experiments, the PRC can be estimated by measuring the phase changes with given stimuli, and several methods have been proposed (reviewed in [3]). One of the problems in such estimation is that the timescales of a PRC, naturally inherited from neuronal dynamics, are around the range of interspike intervals, while the external stimuli, such as pulses or continuous noise, usually have strong power in higher frequency ranges. This problem has been addressed in many ways such as simple averaging [4], randomly selecting low frequency Fourier modes [2,3], assuming smooth priors [5], etc.

Here we propose a systematic and efficient approach of estimating the PRC based on the recently developed method in signal processing called Compressive Sensing (CS) [6]. CS is a framework for estimation of sparsely constructed signals from random observations, and therefore suitable for discovering a small number of low frequency Fourier modes composing the PRC from the experimental measurements with a much wider frequency spectrum.

Using simulated and experimental data, we show that our CS-based method can produce a decent estimate of the PRCs particularly when the number of spikes is so small that averaging cannot help (Fig. 1A, B). Furthermore, since contribution of each mode is systematically evaluated, one can also examine the tradeoffs between the number of modes and goodness-of-fit (Fig. 1C), and




Figure 1 A. PRCs reconstructed from simulated data of the Morris-Lecar model via WSTA [4] and our CS-based method. The number of spikes was 400. In the CS case, the estimated PRCs composed of 200 (blue) and 3 (red) Fourier modes are shown. **B.** Predicted phase shifts vs measured phase shifts in each method. The black line represents the perfect match. **C.** Number of modes vs goodness-of-fit (R^2) in the CS-based method.

* Correspondence: shhong@oist.jp

¹Computational Neuroscience Unit, Okinawa Institute of Science and Technology, Okinawa 904-0411, Japan

Full list of author information is available at the end of the article

therefore pick out most relevant and predictive part of the measurements.

Author details

¹Computational Neuroscience Unit, Okinawa Institute of Science and Technology, Okinawa 904-0411, Japan. ²Theoretical Neurobiology, University of Antwerp, B-2610 Antwerpen, Belgium.

Published: 18 July 2011

References

1. Winfree AT: **The geometry of biological time**. New York: Springer; 2001.
2. Izhikevich EM: **Dynamical systems in neuroscience : the geometry of excitability and bursting**. Cambridge, Mass.: MIT Press; 2007.
3. Torben-Nielsen B, Uusisaari M, Stiefel KM: **A comparison of methods to determine neuronal phase-response curves**. *Front Neuroinformatics* 2010, **4**:6.
4. Ota K, Nomura M, Aoyagi T: **Weighted Spike-Triggered Average of a Fluctuating Stimulus Yielding the Phase Response Curve**. *Phys. Rev. Lett* 2009, **103**:024101.
5. Nakae K, Iba Y, Tsubo Y, Fukai T, Aoyagi T: **Bayesian estimation of phase response curves**. *Neural Netw* 2010, **23**:752-763.
6. Baraniuk R: **Compressive sensing**. *IEEE Signal Processing Magazine* 2007, **24**:118.

doi:10.1186/1471-2202-12-S1-P61

Cite this article as: Hong and De Schutter: Efficient estimation of phase response curves via compressive sensing. *BMC Neuroscience* 2011 **12** (Suppl 1):P61.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

